

# Modeling Wind Wave Evolution from Deep to Shallow Water

Tim T. Janssen

Department of Geosciences, San Francisco State University  
San Francisco, California 94132  
phone: 415 338 1209; fax: 415 338 7705; email: [tjanssen@sfsu.edu](mailto:tjanssen@sfsu.edu)

Thomas H. C. Herbers

Department of Oceanography, Code OC/He, Naval Postgraduate School  
Monterey, California 93943  
phone: 831 6562917; fax: 831 656 2712; email: [thherber@nps.edu](mailto:thherber@nps.edu)

Gerbrant Ph. van Vledder

Department of Environmental Fluid Mechanics, Delft University of Technology  
2600 GA Delft, The Netherlands  
phone: +1 31 15 2781953 ; fax: +1 31 15 2784842 ; email: [g.p.vanvledder@tudelft.nl](mailto:g.p.vanvledder@tudelft.nl)

Award Numbers: N000141010391, N0001410WX10046, N000141010453

## LONG-TERM GOALS

Ocean waves are an important aspect of upper ocean dynamics, in particular on the shallow continental shelves and in coastal areas. The long-term objective of this work is to advance modeling capability in such shallow areas by improving model representations of nonlinear effects and dissipation.

## OBJECTIVES

The specific objectives of the present work are 1) to develop and implement an efficient, scalable approximation for the nonlinear quadruplet source term, 2) to develop and implement a generalized nonlinear source term that is accurate in water of arbitrary depth, 3) to develop and implement an improved nonlinear closure for triad nonlinear interactions in shallow water, and 4) improve representations of dissipation by wave breaking and wave-bottom interactions.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>2010</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2010 to 00-00-2010</b>	
4. TITLE AND SUBTITLE <b>Modeling Wind Wave Evolution from Deep to Shallow Water</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Naval Postgraduate School, Department of Oceanography, Code OC/He, Monterey, CA, 93943</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>6</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

## APPROACH

Modern, third-generation (3G) wave models are based on the action balance (or radiative transfer) equation, which is a geometrical optics description of the evolution of wave energy (or action) through a slowly varying medium and time. In Lagrangian form (for convenience) this balance equation can be written as

$$\frac{dN(\mathbf{k})}{dt} = S_{in}(\mathbf{k}) + S_{ds}(\mathbf{k}) + S_{sc}(\mathbf{k}) + S_{nl}(\mathbf{k}) \quad (1)$$

where  $N(\mathbf{k})$  is the wave action at wavenumber vector  $\mathbf{k}$  and  $t$  is time. The forcing terms on the right-hand side are known as source terms and account for the input of energy by the wind ( $S_{in}$ ), spectral redistribution of energy through scattering by seafloor topography ( $S_{sc}$ ) or through nonlinear wave-wave interactions ( $S_{nl}$ ), and dissipation of wave energy ( $S_{ds}$ ) through e.g. breaking or bottom friction.

In this study we will develop and improve the source terms for nonlinear interactions  $S_{nl}$  and energy dissipation  $S_{ds}$ , to account for effects of finite depth and shallow water, and to ensure a consistent and smooth model representation of wave evolution from deep to shallow water.

### Nonlinearity

We will develop an efficient method for the evaluation of the nonlinear source term, allowing for greater efficiency and accuracy in operational use. To allow modeling of wave propagation from deep to shallow water, we will modify the nonlinear source term to account for changes in relative water depth [Janssen *et al.* 2006], and develop an improved closure approximation for nearshore wave propagation.




Figure 1. Illustration of changes in nonlinear regimes as ocean waves travel across the shelf into shallow water.

### Dissipation

We will develop and test improvements to wave dissipation parameterizations through detailed comparisons against laboratory and field observations.

## WORK COMPLETED

The completed work in this project thus far includes:

- Analysis of dissipation in one-dimensional nonlinear shoaling waves (laboratory data set by Boers, 1996) using bispectral analysis.
- Development and preliminary testing of nonlinear shallow-water wave model using a new closure approximation.
- Development and preliminary testing of a new scaling for the breaking parameter  $\gamma$ .
- Completed review of formulations for nonlinear four-wave interactions in shallow water.

## RESULTS

### Dissipation in nonlinear shoaling waves

Through bispectral analysis of laboratory observations (reported in Boers 1996) we have analyzed the dissipation spectrum in breaking waves. Our findings confirm that in the very nearshore the dissipation is approximately weighted as  $f^2$  (frequency-squared) and the dissipation spectrum approaches a white-noise signature [e.g. Kaihatu et al. 2007]. Further, we find that the weighting of dissipation varies across the flume (figure 2) and for varying incident wave conditions. This dependency is being further investigated.




Figure 2. Frequency exponent values from regression analysis for dissipation spectrum and flux spectrum tail as a function of relative water depth.

### A nonlinear model for shallow-water wave propagation

Using the results from the dissipation analysis, and implementing an empirical closure approximation, a research model is developed to test the principles and ideas underlying our approach. Model results are in good agreement with flume observations of spectra and higher-order bulk statistics such as skewness and asymmetry (figure 3). These preliminary results show the potential of our modeling approach. A more rigorous modeling effort, borrowing ideas from turbulence renormalized perturbation [e.g. McComb, 1990] theory but using the same basic principles, is presently under development.




Figure 3. Evolution of asymmetry, a third-order bulk statistics, for Boers experiment 1C (Boers 1996).

### Development of a new scaling of breaking parameter $\gamma$

The newly developed scaling of the breaking parameter  $\gamma$ , referred to here as  $n$ -kd scaling, is implemented, calibrated and tested against field and laboratory observations. Comparisons between model predictions and laboratory observations for both the traditional fixed-value  $\gamma$  and the new  $n$ -kd scaling (calibrated with field and laboratory observations) show improved model performance (figure 4). Further testing of the new scaling is presently ongoing [Holthuijsen & Salmon, 2010].




Figure 4 Scatter plots for predicted and observed significant wave height for a number of laboratory studies.

Left panel: breaker index default fixed value  $\gamma = 0.73$  . Right panel: new  $n - kd$  scaling for  $\gamma$  .

## IMPACT/APPLICATIONS

### Economic Development

The improvements to coastal wave prediction models as developed in this project, will contribute to various industries operating on the continental shelf and the coastal zone, such as fisheries, and shipping and offshore industry. Further, the availability of better wave prediction models will benefit coastal and ocean engineering companies e.g. in the design and operation of offshore and coastal structures, and the development of coastal management strategies.

### Quality of life

The improvements to coastal wave prediction models as developed in this project, will improve modeling capability of coastal circulation and transport processes, which will benefit coastal recreation (more reliable knowledge of wave heights, rip currents etc), coastal management, and help mitigate pollution hazards for humans (recreation) and coastal ecosystems.

## TRANSITIONS

### Economic Development

The developments in this project will be made available as open source software and as modules to widely used operational wave models. These models are used by NOAA and other agencies involved in coastal development and management, and by many coastal and ocean engineering companies.

### Quality of life

The software developed within this project will be disseminated in open source models used by local and federal agencies and companies involved in coastal recreation (surf prediction, rip currents, pollution), coastal management, and mitigation of coastal hazards.

## REFERENCES

Boers, M. (1996) Simulation of a surf zone with a barred beach, part 1: Wave heights and wave breaking, *Comm. Hydr. Geotechn. Engn.* 96-5 Delft University of Technology, 116 pp.

Holthuijsen, L. H. and J. Salmon (2010) The n-kd scaling of depth-induced wave breaking. *manuscript in preparation*.

Janssen, T. T., T.H.C. Herbers, and J. A. Battjes (2006) Generalized evolution equations for nonlinear surface gravity waves over two-dimensional topography, *J. Fluid Mech.* **552**, 393-418.

Kaihatu, J. M., Veeramony. J., Edwards, K. L., and Kirby, J. T. (2007) Asymptotic behavior of frequency and wavenumber spectra of nearshore shoaling and breaking waves. *J. Geoph. Res.* **112**, doi:10.1029/2006JC003817.

McComb, W. D. (1990) *The physics of fluid turbulence* Clarendon Press, Oxford, 572 pp.